

Domination and Disappearance of the Black Sponge: A Quarter Century after the Initial *Terpios* Outbreak in Southern Japan

James D. Reimer^{1,*}, Yoko Nozawa², and Euichi Hirose³

¹Rising Star Program, Univ. of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan
²Biodiversity Research Center, Academia Sinica, Taipei 115, Taiwan
³Department of Chemistry, Biology and Marine Science, Faculty of Science, Univ. of the Ryukyus, 1 Senbaru, Nishihara, Okinawa 903-0213, Japan

(Accepted May 7, 2010)

The encrusting cyanobacteriosponge *Terpios hoshinota*, reported from western-Pacific coral reefs, competes for space by encrusting reef substrata including live corals, and has been expanding its range in the northwestern Pacific Ocean (Liao et al. 2007). This species occasionally undergoes massive outbreaks which can cause considerable damage to reefs (Bryan 1973, Rützler and Muzik 1993). In Japan, the 1st outbreak was reported in 1986 at Yonama, Tokunoshima I. (Fig. 1), with up to 87.9% cover, killing most corals in the area (Marine Park Center Foundation 1986). Since then; however, very little research has been conducted on *T. hoshinota*, and the nature and causes of the outbreaks remain unknown. In early spring 2010, Tokunoshima I. and neighboring Okinoerabu I. were investigated to determine the



Fig. 1. Map showing locations investigated in this study. (A) Location of Tokunoshima and Okinoerabu. in the western Pacific Ocean. (B) Locations of the past outbreak (Yonama) and current outbreak (Yakomo) of *Terpios hoshinota*.

current status of *T. hoshinota*. Surprisingly, no *T. hoshinota* patches were present at Yonama, and much of the site was covered in dense *Acropora* colonies (Fig. 2). Furthermore, while there was little or no *T. hoshinota* present at most other examined sites on both islands, a new massive outbreak was confirmed at Yakomo, Okinoerabu I. (Fig. 1), with total cover often exceeding 50% according to transect data (Figs. 3, 4). These preliminary results suggest that *T. hoshinota* outbreaks, although very destructive, may be sporadic in nature, and not always an irreversible "climax" for affected reefs. However, the continued presence of *T. hoshinota* almost 25 yr after its initial outbreak indicates that this species is able to persist once it enters a reef. http://zoolstud.sinica.edu.tw/Journals/50.3/394.pdf



Fig. 2. Dense Acropora colonies with no Terpios hoshinota at Yonama.



Fig. 3. Example of high coverage of *Terpios hoshinota* in a transect at Yakomo.



Fig. 4. *Terpios hoshinota* overgrowing live coral at Yakomo.

Acknowledgments: We appreciate funding from the Ocean Exposition Commemorative Park Management Foundation, Motobu, Okinawa, Japan.

REFERENCES

Bryan P. 1973. Growth rate, toxicity, and distribution of the encrusting sponge, *Terpios* sp. (Hadromerida: Suberitidae) in Guam, Mariana Islands. Micronesica **9:** 237-242.

Liao M, S Tang, C Hsu, K Wen, W Chen, J Wang et al. 2007. The "black disease" of reef-building corals at Green Island, Taiwan - outbreak of a cyanobacteriosponge, *Terpios hoshinota* (Suberitidae: Hadromerida). Zool. Stud. **46:** 520.

Rützler K, K Muzik. 1993. Terpios hoshinota, a new cyanobac teriosponge threatening Pacific reefs. Sci. Mar. 57: 395-403.

Marine Park Center Foundation. 1986. Urgent survey report of an abnormal phenomenon of marine ecosystem in the Amami Islands. Tokyo: Environmental Agency of Japan, 39 pp. (in Japanese)

^{*}To whom correspondence and reprint requests should be addressed. Tel: 81-98-8958542. Fax: 81-98-8958576. E-mail:jreimer@sci.u-ryukyu.ac.jp