

- Holm, T. 1904. Studies upon the Cyperaceae 23. The inflorescence of Cyperus in North America. Am. J. Sci. 18:301-7.
- Holttum, R. E. 1948. The spikelet in Cyperaceae. Bot. Rev. 14:525-41.
- Koyama, T. 1961. Classification of the family Cyperaceae (1). J. Fac. Sci. Tokyo Univ. (Bot.) 8:37-148.
- Kuoh C. S. 1974. A Systematic study on the Formosan Cyperaceae (1) Morphological study of the floral parts. Biol. Bull. Taiwan normal univ. no. 9. 47-57.
- 1975. On the distribution of Formosan Carex (Cyperaceae). ibid. no. 10. 43-52.
- Ohwi, J. 1944. Cyperaceae Japonicae. Pt. 2. Mem. Coll. Sci. Kyoto Imp. Univ. B, 18:1-182.
- Shah, C. K. 1962. Floral morphology of the Cyperaceae. I. Naturaliste Can. 89:330-44.

Abstract

In this study, the author recognize 25 taxa of the genus Cyperus, in which C. pannonicus (= Juncellus pannonicus) has first been reported from Formosa. As regards to the arrangement of the taxa, the systematic information such as prickly hairs of leaves, types of inflorescence, floral scales, intra-spicular bracts, prophylls and extra-spicular prophylls have been taken into consideration. Attention has been paid especially to the characters of the intra-spicular prophylls which is the diagnostic characters in this genus.

智慧藏