

with red margins along anther sacs (Figure 2:H). A comparison of the salient characters of the four species in sect. *Coelocentrum* is shown in Table 1.

Acknowledgments. We thank Hidetoshi Nagamasu (KYO) for assistance with the Latin diagnosis; Hong-Zhe Li and Hong Ma (KUN) for providing handsome images (Figure 8: B, D) and useful literature; Dave Boufford (A/GH) and Hidetoshi Nagamasu for improving the manuscript; and the curators of GXMI, K and KUN for facilitating our examination of herbarium collections. This study was supported in part by the Guangxi Natural Science Foundation (GKJ0448089) to Yan Liu (IBK) and grants from the Research Center for Biodiversity, Academia Sinica and the National Science Council, Taiwan, to Ching-I Peng (HAST).

Literature Cited

- Ku, S.M., C.-I Peng, and Y. Liu. 2004. Notes on *Begonia* (sect. *Coelocentrum*, Begoniaceae) from Guangxi, China, with the report of two new species. Bot. Bull. Acad. Sin. **45**: 353-367.
- Legro, R.A.H. and J. Doorenbos. 1969. Chromosome numbers in *Begonia* 1. Netherlands J. Agric. Sci. **17**: 189-202.
- Peng, C.-I, Y.M. Shui, Y. Liu, and S.M. Ku. 2005a. *Begonia fangii* (sect. *Coelocentrum*, Begoniaceae), a new species from limestone areas in Guangxi, China. Bot. Bull. Acad. Sin. **46**: 83-89.
- Peng, C.-I, S. M. Ku, and W.C. Leong. 2005b. *Begonia liuyanii* (sect. *Coelocentrum*, Begoniaceae), a new species from limestone areas in Guangxi, China. Bot. Bull. Acad. Sin. **46**: 245-254.
- Peng, C.-I, Y. K. Chen, and W.C. Leong. 2005c. Five new species of *Begonia* (Begoniaceae) from Taiwan. Bot. Bull. Acad. Sin. **46**: 255-272.
- Shui, Y.M. and W.H. Chen. 2005. New data of sect. *Coelocentrum* (*Begonia*) in Begoniaceae. Acta Bot. Yunnan. **27**(4): 355-374.
- Shui, Y.M., C.-I Peng, and C.Y. Wu. 2002. Synopsis of the Chinese species of *Begonia* (Begoniaceae), with a reappraisal of the sectional delimitation. Bot. Bull. Acad. Sin. **43**: 313-327.
- Ye, H.G., F.G. Wang, Y.S. Ye, and C.-I Peng. 2004. *Begonia coptidifolia* (Begoniaceae), a new species from China. Bot. Bull. Acad. Sin. **45**: 259-266.

中國廣西石灰岩地區秋海棠屬側膜組一新種：一口血秋海棠

劉 演¹ 古訓銘^{2,3} 彭鏡毅²

¹ 廣西壯族自治區 中國科學院廣西植物研究所

² 中央研究院生物多樣性研究中心 植物標本館

³ 國立成功大學生命科學系

本文報導中國廣西壯族自治區西南部石灰岩地區的秋海棠屬側膜組 (*Begonia* sect. *Coelocentrum*) 的一個新種：一口血秋海棠 (*Begonia picturata*)，提供線繪圖與彩色照片以資辨識，並報導其染色體數 ($2n = 30$)。一口血秋海棠的花梗具長曲柔毛，小花梗、外花被片、子房與果實具有紅色圓錐狀的彎曲剛毛或直硬毛而與同為側膜組的捲毛秋海棠 (*B. cirrosa*) 略似；但一口血秋海棠葉片具有花紋、上表面密布長曲柔毛狀剛毛或絨毛狀剛毛，捲毛秋海棠葉片無斑紋、上表面被疏至中度的柔毛或剛毛狀柔毛。本文並提供同為側膜組較相似之龍州秋海棠 (*B. morsei*) 之模式標本與彩紋秋海棠 (*B. variegata*) 的彩色照片，以資辨識。

關鍵詞：捲毛秋海棠；彩紋秋海棠；龍州秋海棠；一口血秋海棠；側膜組；秋海棠科；中國；石灰岩植物；廣西；染色體數；藥用植物；新種；稀有植物。